


BYSTANDER AED USE

STATS YOU MAY NOT KNOW



90%

of victims who receive a shock from an AED in the first minute **live.**¹

The chances of survival decrease by

7-10%

for every minute that passes without CPR and defibrillation.²



Victims have

2.7x



higher odds of survival to hospital discharge, with good neurological outcomes, when they receive a shock from a bystander than victims who have to wait for EMS arrival for a shock.³

70%

of cardiac arrest patients who don't receive AED therapy from a bystander will either not survive or survive with impaired brain function.⁵


60%



of public out-of-hospital cardiac arrests have shockable rhythms.⁴

3,459

lives could be saved (with good neurological outcome!) annually in the U.S., if 100% of bystanders apply an AED.⁶



BUT...



Only 11%

of out-of-hospital cardiac arrest victims have an AED applied by a bystander.⁷



63%

of Americans report a lack of confidence in using an AED.⁸



2%

of bystanders are willing to use an AED.⁹

Reasons include:

- Not knowing how the AED works
- Fear of causing harm to the patient
- Not being comfortable using the AED
- Legal liability

Empowering bystanders to use AEDs is crucial to improving outcomes from sudden cardiac arrest. The latest training technology can make it easier for you to ensure your learners are mastering this critical link in the chain of survival.

Learn more at Laerdal.com

Sources:
 1. AED Fact Sheet. (n.d.) American Heart Association. Retrieved from <https://cpr.heart.org/-/media/CPR-Files/Training-Programs/AED-Implementation/2023-updates/KJ1683-AED-Program-Checklist.pdf>
 2. Implementing an AED Program. (2018). American Heart Association. Retrieved from <https://cpr.heart.org/-/media/CPR-Files/Training-Programs/AED-Implementation/2022-updates/AED-Guide-1.pdf>
 3. Pollack, R. A., Brown, S. P., Rea, T., Aufderheide, T., Barbic, D., Buick, J. E., Christenson, J., Idris, A. H., Jasti, J., Kampp, M., Kudenchuk, P., May, S., Muhr, M., Nichol, G., Ornato, J. P., Sopko, G., Vaillancourt, C., Morrison, L., & Weisfeldt, M. (2018). Impact of bystander automated external defibrillator use on survival and functional outcomes in shockable observed public cardiac arrests. *Circulation*, 137(20), 2104–2113. <https://doi.org/10.1161/circulationaha.117.030700>
 4. Ibid.
 5. Study Reports Cardiac Arrest Survival Greatly Increases When Bystanders Use an AED. (2018). *Journal of Emergency Medical Services*. (2018). Retrieved from <https://www.jems.com/news/study-reports-cardiac-arrest-survival-greatly-increases-when-bystanders-use-an-aed/>
 6. Pollack, R. A., Brown, S. P., Rea, T., Aufderheide, T., Barbic, D., Buick, J. E., Christenson, J., Idris, A. H., Jasti, J., Kampp, M., Kudenchuk, P., May, S., Muhr, M., Nichol, G., Ornato, J. P., Sopko, G., Vaillancourt, C., Morrison, L., & Weisfeldt, M. (2018). See reference #3.
 7. CARES Annual Report: 2022. Cardiac Registry to Enhance Survival (CARES). Retrieved from https://mycares.net/stepages/uploads/2023/2022_flipbook/index.html?page=2
 8. Survey Reveals Nearly Two Thirds of Americans are Not Confident They Know How to Operate an AED. (2019). Sudden Cardiac Arrest Foundation. Retrieved from <https://www.sca-aware.org/sca-news/survey-reveals-nearly-two-thirds-of-americans-are-not-confident-they-know-how-to-operate-an-aed>
 9. Smith, C. M., Lim, Choi, Kaung, S. N., Khan, M. O., Arvanitis, T. N., Fothergill, R., Hartley-Sharp, C., Wilson, M. H., & Perkins, G. D. (2017). Barriers and facilitators to public access defibrillation in out-of-hospital cardiac arrest: A systematic review. *European Heart Journal - Quality of Care and Clinical Outcomes*, 3(4), 264–273. <https://doi.org/10.1093/ehjqcco/qcx023>

©2023 Laerdal Medical. All rights reserved. 2308-1182882828